

The interplay of soft and hard contributions in the electromagnetic pion form factor¹

W. Schweiger^a

^aInstitut für Theoretische Physik, Karl-Franzens-Universität Graz, A-8010 Graz, Austria

We consider various relativistic models for the valence Fock-state wave function of the pion. These models are obtained from simple instant-form wave functions by applying a Melosh rotation to the spin part and by imposing physical constraints on the parameters. We discuss how the soft and the hard (perturbative) parts of the electromagnetic form factor are affected by the choice of the model and by the Melosh rotation.

The knowledge of wave functions of the hadronic constituents allows to link hadronic phenomena in different kinematical regions. For several reasons it is advantageous to consider the composition of hadrons out of its constituents at fixed light-front time $\tau = t + z$ rather than at ordinary time t . The “time” evolution in τ is then determined by front-form dynamics. One of the attractive features of such an approach is that the corresponding wave functions are direct generalizations of non-relativistic wave functions in the sense that they can be interpreted as probability amplitudes for finding a particular Fock state in the hadron under consideration with the constituents carrying certain momenta, spins, etc. Like in the non-relativistic case light-front wave functions can be expressed in terms of purely internal variables (momentum fractions x and transverse momenta $k_\perp$).

For a hadron form factor the, in principle, exact expression is just a sum (over all Fock states) of overlap integrals of incoming and outgoing light-front wave functions [1]. A widely used approximation is then to assume that the dominant contribution comes from the valence Fock state. For large momentum transfers $Q \rightarrow \infty$ the analysis of the corresponding overlap integral reveals that the one-gluon-exchange tail of the wave function can be factored out, so that one ends up with a perturbative representation of the form factor in terms of a convolution integral [2]. The distribution amplitude $\phi(x, \tilde{Q})$ entering this convo-

lution integral is again related to the valence-quark light-front wave function of the hadron. Its dependence on the factorization scale $\tilde{Q}$ (which in turn depends on Q) is given by an evolution equation which is driven by one-gluon-exchange. Practically, this means that the knowledge of the soft part of the wave function suffices, since the high-momentum tail of the wave function is determined by perturbative evolution. What we therefore want to model is only the soft part of the pion wave function.

A commonly used ansatz for the quark-antiquark light-front wave function of the pion is of harmonic-oscillator type

$$\psi(x, k_\perp) = A_\pi \chi(x, k_\perp) J(x, k_\perp) \exp\left(-\frac{M_0^2}{8\beta^2}\right) \quad (1)$$

with M_0^2 denoting the front-form expression for the free two-particle mass

$$M_0^2 = \frac{k_\perp^2 + m_q^2}{x(1-x)}. \quad (2)$$

A_π is a normalization constant, $\chi(x, k_\perp)$ the (light-front) spin wave function of the $q\bar{q}$ pair, and $J(x, k_\perp)$ the square root of a Jacobian. Such a model for the pion wave function has, e.g., been proposed by Brodsky, Huang and Lepage (BHL) [3]. They took $J = 1$ and the usual (instant-form) expression for the spin-wave function χ . Terent'ev and Karmanov (TK) [4], on the other hand, chose $J(x) = \sqrt{1/2x(1-x)}$, i.e. the square root of the Jacobian relating

¹Talk given at the “International Light-Cone Workshop”, Trento, Italy, September 2001

Table 1

Parameters and properties of the wave-function models considered.

wave function	β (MeV)	A_π (GeV ⁻¹)	$P_{q\bar{q}}$	$\sqrt{\langle k_\perp^2 \rangle}$ (MeV)
ψ_{BHL} (full MR)	300	110.	1.00	282
ψ_{BHL} (no MR)	300	78.2	0.51	282
ψ_{BHL} (approx. MR [11])	300	65.8	1.85	323
ψ_{TK} (full MR)	291	77.5	1.00	270
ψ_{CCP} (full MR)	272	80.0	1.00	272
ψ_{PL} (full MR)	1045	238.	1.00	266

 $m_q = 330$ MeV and $f_\pi = 93$ MeV throughout.

the relativistic integration measures (d^3k/k^0) and ($d^2k_\perp dx$). The $J(x, k_\perp)$ adopted by Chung, Coester and Polyzou (CCP) [5] was of the form $J(x, k_\perp) = \sqrt{M_0/4m_q x(1-x)}$, i.e. the square root of the Jacobian relating the non-relativistic integration measure (d^3k/m_q) to ($d^2k_\perp dx$). In addition to these different choices of the J s we will use another simple ansatz for the s-state orbital function of the pion, which has a power-law form

$$\psi_{\text{PL}}(x, k_\perp) = A_\pi \chi(x, k_\perp) J(x, k_\perp) \left(\frac{\beta^2}{M_0^2 + \beta^2} \right)^\alpha.$$

A few words about the spin wave function $\chi(x, k_\perp)$ are also in order. It is a well known disadvantage of the light-front formalism that the usual addition of angular momenta (e.g. for 2 particles $\vec{J} = \vec{L} + \vec{S}^{(1)} + \vec{S}^{(2)}$) holds only for the third component, whereas the addition law for the other two components is, in general, much more complicated. It is, however, possible to go over to a unitarily transformed spin operator which satisfies the usual spin-addition laws [4]. This unitary transformation is known as the ‘‘Melosh rotation’’ (MR). The light-front spin wave function $\chi(x, k_\perp)$ results thus from an inverse Melosh rotation of an ordinary spin wave function. A convenient expression for $\chi(x, k_\perp)$ can, e.g., be found in Ref. [6]. Remarkably, the light-front spin wave function of an s-wave pion contains also helicity ± 1 components.

The idea behind the construction of such wave-function models is always that the valence-Fock-state wave function within QCD has something to do with the wave function of constituent

quarks (which can be obtained from much simpler dynamics). Therefore the mass m_q occurring in the wave functions has to be interpreted as constituent-quark mass. We took $m_q = 330$ MeV. According to the finding in Ref. [7] we have also fixed the parameter α in ψ_{PL} to be $\alpha = 3.5$. The normalization A_π has then been adjusted such that the weak decay constant $f_\pi = 2\sqrt{3} \int dx d^2k_\perp \psi = 93$ MeV is reproduced. Finally, the parameter β has been determined (for the full Melosh rotated wave functions) by means of the valence-quark dominance assumption $P_{q\bar{q}} = \int dx d^2k_\perp \psi^* \psi = 1$. This also lead to reasonable values for the mean intrinsic transverse momenta $\sqrt{\langle k_\perp^2 \rangle}$ of a quark (or an antiquark) inside the pion (cf. Table 1 where the model parameters are also summarized).

The dependence of the soft (overlap) contribution to F_π on the choice of the quark-antiquark wave function is displayed in Fig. 1. At very small Q^2 (small figure) the curves nearly coincide. Figure 2, on the other hand, shows the interplay of soft and hard (perturbative) contributions to F_π and the influence of the MR factor. Results are only considered for ψ_{BHL} , since the hard contribution to the form factor is nearly independent of the choice of the wave function. It does not even depend on whether the MR factor is fully taken into account or neglected at all. Marked differences are only observed if the MR factor is approximated like in Ref. [11]. Shortly summarized, the following conclusions can be drawn from the figures (and from Table 1):

The use of different wave functions does not have a significant effect on the electromagnetic pion




Figure 1. Soft contributions to $Q^2 F_\pi$ (large figure) and to F_π^2 (small figure) in different Q^2 ranges for ψ_{BHL} (solid line), ψ_{TK} (short-dashed line), ψ_{CCP} (medium-dashed line), ψ_{PL} (long-dashed line) with the full MR factor and for ψ_{BHL} without MR factor (dotted line). Data are taken from Ref. [8] (filled triangles), Ref. [9] (filled circles), and Ref. [10] (open circles).

form factor as long as these wave functions provide comparable results for f_π and $\langle k_\perp^2 \rangle$. This holds in both, the soft and the hard regimes.

Normalizing the wave functions in such a way that f_π remains unaltered, the soft part of F_π (and also $P_{q\bar{q}}$) becomes considerably smaller if the MR factor is neglected. Minor changes due to the omission of the MR factor can be observed in the perturbative part of F_π . The approximations for the MR factor adopted in Ref. [11], on the other hand, entail also sizable changes in the hard part of F_π .

Comparing the soft with the perturbative contribution to F_π one observes that the latter starts to dominate at $Q^2 \approx 5 \text{ GeV}^2$.

The effect of the MR factor on the pion distribution amplitude is just a slight broadening as compared to the asymptotic distribution amplitude (cf. Fig. 2). The broadening becomes much stronger if the approximations of Ref. [11] are applied to the MR factor. A recent QCD sum-rule analysis of the pion distribution amplitude by Bakulev et al. [12] provides a double-humped distribution amplitude (cf. Fig. 2), but with less pronounced structure than originally proposed by Chernyak and Zhitnitsky.




Figure 2. Soft (solid line) and hard (dash-dotted line) contributions to $Q^2 F_\pi$ for ψ_{BHL} with the full MR factor. Results without (dotted line) and with approximate treatment [11] (dash double-dotted lines) of the MR factor are shown for comparison. Corresponding pion distribution amplitudes are displayed in the small figure along with a recently proposed model [12] (grey line).

REFERENCES

1. S.D. Drell and T.M. Yan, Phys. Rev. Lett. 24 (1970) 181.
2. G.P. Lepage and S.J. Brodsky, Phys. Rev. D22 (1980) 2157.
3. S.J. Brodsky, T. Huang and G.P. Lepage, in Particles and Fields, eds. A.Z. Capri and A.N. Kamal (Plenum Press, New York, 1983).
4. M.V. Terent'ev, Sov. J. Nucl. Phys. 24 (1976) 106; V.A. Karmanov, Nucl. Phys. B166 (1980) 378.
5. P.L. Chung, F. Coester and W.N. Polyzou, Phys. Lett. B205 (1988) 545.
6. B.-Q. Ma, Z. Phys. A345 (1993) 321.
7. F. Schlumpf, Phys. Rev. D47 (1993) 4114.
8. C.J. Bebek et al., Phys. Rev. D17 (1978) 1693.
9. J. Volmer et al., Phys. Rev. Lett. 86 (2001) 1713.
10. S.R. Amendolia et al., Nucl. Phys. B277 (1986) 168.
11. Z. Dziembowski and L. Mankiewicz, Phys. Rev. Lett. 21 (1987) 2175.
12. A.P. Bakulev, S.V. Mikhailov and N.G. Stefanis, Phys. Lett. B508 (2001) 279.